PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:		(11) International Publication Number:	WO 94/22496
A61K 49/02, 43/00	A1	(43) International Publication Date:	13 October 1994 (13.10.94)
(21) International Application Number: PCT/US (22) International Filing Date: 29 March 1994 (2)		PL, SK, European patent (AT,	BE, CH, DE, DK, ES, FR,
(30) Priority Data: 08/040,739 31 March 1993 (31.03.93)	τ	Published With international search repo	n.
(71) Applicant: MALLINCKRODT MEDICAL, INC. 675 McDonnell Boulevard, P.O. Box 5840, St. Lo. 63134 (US).			
(72) Inventors: BRODACK, James, W.; 3827 Kentuck Drive, Florissant, MO 63034 (US). DEROSCH, M 1762 Winding Glen Drive, St. Charles, MO 633 DEUTSCH, Edward, A.; 12805 Maryland Estate Maryland Heights, MO 63043 (US). DEUTSCH F.; 12805 Maryland Estates Court, Maryland Heights, MO 63043 (US). DYSZLEWSKI, Mary, Marmion Autumn Dale Court, Maryland Heights, MO 63043 (74) Agents: MCBRIDE, Thomas, P. et al.; Mallinckrodt	Mark, 1 03 (US es Cou I, Kare ghts, M i; 122: 3 (US)	A.; S). rt, m, iO O 21	
Inc., 675 McDonnell Boulevard, P.O. Box 5840, S MO 63134 (US).			

(54) Title: RADIOPHARMACEUTICAL FORMULATIONS HAVING NON-STANNOUS REDUCTANTS

(57) Abstract

The present invention relates to novel radiopharmaceutical imaging agents having non-stannous reductants. Metallic compounds, such as Cu(I), Cu(II), Co(II), Fe(II), Sn(O), Zr(O), Cr(II) and Zn(O), will act to effectively reduce radionuclide containing solutions. Several non-metallic compounds, such as acids in general, dithionite, formamidine, formamidine sulfinic acid, phosphite, hypophosphite, dithiotreitol, hydrochloric acid, and borohydric acid may also be used to reduce radionuclide containing solutions. Moreover, it has been discovered that several agents, such as phosphines sulfhydryl compounds, phosphites, thiols, thioethers, borates, borocyano groups, ascorbates and gentisates efficiently reduce radionuclide containing solutions and complex with the radionuclide at the same time. The present invention also relates to kits for forming radiopharmaceutical imaging agents, such kits including non-stannous reducing agents.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MIR	Mauritania
ΑŪ	Australia	GB	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	Œ	Ireland	NZ	New Zealand
BJ	Benin	π	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KR	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF.	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CG	Congo		of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SI	Slovenia
α	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovakia
CM	Cameroon	Ц	Liechtenstein	SN	Scoegal
CN	China	LK	Sri Lanka	TD	Ched
CS	Czechoslovakia	LU	Luxenbourg	TG	Togo
CZ	Czech Republic	LV	Letvia	Tj	Tajikistan
DR	Germany	MC	Mozaco	77	Trimidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
Fī	Finland	MIL	Mali	UZ	Uzbekistan
FR	France	MIN	Mongolia	VN	Viet Nam
GA	Gabon		•		

WO 94/22496 PCT/US94/03389

RADIOPHARMACEUTICAL FORMULATIONS HAVING NON-STANNOUS REDUCTANTS

Background

The present invention relates to novel radiopharmaceutical imaging agents having non-stannous reductants. The present invention further relates to kits for forming radiopharmaceutical imaging agents, such kits including non-stannous reducing agents.

5

10

15

20

25

30

Several non-invasive methods of imaging skeletal structures, and body organs and tissues have been developed over the past decades. These methods are based on the tendency of the particular skeleton, organ or tissue to concentrate certain chemicals which may be detectable, such as through the use of scintiphotography or radiation detection. The use of radiopharmaceutical imaging agents in imaging skeletal structures, organs and tissues, is well known in the fields of biological and medical research as well as diagnostic evaluation procedures. Metal-based radiopharmaceuticals, such as those based on technetium have been found to provide particularly useful images of skeletal structures, and body organs and tissues from which diagnostic information may be obtained. More particularly, radiopharmaceuticals based on technetium 99m have been used successfully as diagnostic imaging agents.

In addition, metal-based radiopharmaceuticals, such as those based on rhenium have been found to be useful as therapeutic agents in the treatment of various diseases. More particularly, radiopharmaceuticals based on rhenium 186 or rhenium 188 have been used successfully as therapeutic agents.

The radiopharmaceutical agents generally include a metal radionuclide, various ligands for binding the

radionuclide to the desired skeletal structure, organ or tissue, reducing agents, stabilizing agents, carriers and delivery vehicles suitable for injection into, or aspiration by a patient, etc.

5

10

15

Because of the relatively short half-lives of the metal radionuclide used in the radiopharmaceutical agent, it is desirable to provide the non-radioactive components of the agent as a kit to which a radionuclide containing solution may be added to form the agent. In particular, a radionuclide generator may be employed in a known manner to obtain a radionuclide which may then be combined and reacted with the contents of a kit which contains appropriate radiopharmaceutical forming components. For example, when forming a technetium imaging agent, a pertechnetate solution may be obtained from a technetium generator. The pertechnetate solution may then be combined and reacted with the components of a kit containing the other materials and agents necessary for forming the radiopharmaceutical agent.

20

25

30

A reducing agent is a necessary component of many radiopharmaceutical kits, the reducing agent acting to reduce the radionuclide containing solution, such as a final the obtain solution, to pertechnetate A reducing agent must be radiopharmaceutical agent. formation of technetium for the in kits radiopharmaceuticals. Stannous ion is the most widely used for forming metal-based kits in reducing agent radiopharmaceuticals. This includes known kits for forming technetium 99m diagnostic agents for imaging the heart, kidney, lungs, and hepatobiliary system, as well as kits for imaging and therapeutic treatment of the brain and skeleton. However, the use of stannous ion as a reducing agent has several disadvantages generally arising from the

5

15

20

25

inherent problems related to the complicated solid and solution chemistry of stannous compounds. In particular, the stannous ion is often a "non-innocent" reducing agent which interferes with or is incorporated into the final radiopharmaceutical.

Therefore it is desirable to provide radiopharmaceutical forming kits which contain non-stannous reducing agents.

Objects Of The Invention

It is one object of the present invention to provide radiopharmaceutical formulations having non-stannous reductants.

It is another object of the present invention to provide kits for forming radiopharmaceutical agents, such kits including non-stannous reducing agents.

Summary Of The Invention

The above objects and others are achieved by providing a kit for forming radiopharmaceutical agents, wherein the kits include non-stannous reducing agents.

Detailed Description Of The Invention

The use of stannous ion as a reductant for preparation of radiopharmaceutical agents is well known. Therefore, stannous ion is commonly included as a component of kits for forming radiopharmaceutical agents, such as agents for imaging the heart, kidneys, lungs and hepatobiliary system and agents for imaging and therapeutic treatment of the

WO 94/22496

5

10

15

20

25

30

Λ

brain and skeleton.

However, the use of stannous ion as a reducing agent in the formation of radiopharmaceutical agents has several disadvantages. In particular, stannous compounds have inherent disadvantages associated with their complicated solid and solution chemistry. For example, stannous ion often acts as a "non-innocent" reductant and may interfere with the formation of the final radiopharmaceutical agent, or become incorporated into the final radiopharmaceutical agent.

It has been discovered that there are a number of pharmaceutically acceptable non-stannous reducing agents which do not possess the same disadvantages of using stannous ion as a reducing agent in the formation of radiopharmaceutical agents.

In particular, metallic compounds, such as Cu(I), Cu(II), Co(II), Fe(II), Sn(0), Zr(0), Cr(II) amd Zn(0), will act to effectively reduce a radionuclide containing solutions, such as pertechnetate solutions, to obtain the desired final radiopharmaceutical agent.

Further, several non-metallic compounds, such as acids in general, dithionite, formamidine, formamadine sulfinic acid, phosphite, hypophosphite, dithiothreitol, hydrochloric acid, and borohydric acid, may also be effectively used to reduce radionuclide containing solutions.

Moreover, it has been discovered that several agents, such as phosphines, sulfhydryl compounds, phosphites, thiols, thioethers, borates, borocyano groups, ascorbates, and gentisates, efficiently reduce the radionuclide

5

10

15

20

25

30

containing solution and complex with the radionuclide at the same time. This is very advantageous in reducing the number of components that must be included in kits for forming radiopharmaceuticals, and in simplifying the chemistry needed to produce the final radiopharmaceutical agent.

According to one embodiment of the present invention, a kit for forming a technetium myocardial imaging agent includes tris(3-methoxypropyl)phosphine (TMPP), as both a reducing agent and as a complexing agent. Further components, such as cuprous ascorbate, may also be included in the kit to increase radiopharmaceutical yield. Notably, it has been found that the addition of stannous ion to the kit actually reduces the product yield by forming reduced hydrolyzed technetium as a by-product.

In a further embodiment of the present invention, a forming radiopharmaceutical imaging includes tertiary phosphines (PR3) wherein the phosphine also acts as a ligand for the technetium complex. particular, it has been discovered that "99mTc(VII)O4 may be reduced using a monodentate phosphine, such as, tertiary phosphines in the presence of a Schiff base ligand (L4). The kit may be a lyophilized kit containing the tertiary phosphine, the Schiff base ligand, and a buffer, but does not require ancillary reductants such as stannous ion, hypophosphite, or ascorbate to carry out the reduction if the technetium-99m generator eluant reaction, It is believed that the degassed prior to its use. reduction reaction proceeds through a 99mTc(V) intermediate, ultimately $^{99m}Tc(V)(0)(L4)^{+}$ to 99mTc(III)(L4)(PR3)2, as a myocardial imaging agent.

In addition, in accordance with the present invention,

5

10

15

20

it has been discovered that hypophosphite ion (H₂PO₂) may be used as an antioxidant and reductant in the formation of Tc-99m radiopharmaceuticals. The reduction potential for hypophosphite ion is comparable to ascorbic acid. Further, the oxidation product of hypophosphite is phosphite, PO₃³, which is totally innocuous in Tc-99m radiopharmaceutical preparations. Hypophosphite and phosphite are totally colorless, easy to analyze, lyophilizable, and injectable.

In another embodiment according to the present invention, it has been discovered that hydrogen phosphite (HPO₃²⁻) may be used as an antioxidant and reductant in the formation of Tc-99m radiopharmaceuticals. The reduction potential for hydrogen phosphite is comparable to ascorbic acid. Further, the oxidation product of hydrogen phosphite is phosphate, PO₄³⁻, which is totally innocuous in Tc-99m radiopharmaceutical preparations. Phosphite and phosphate are colorless, easy to analyze, lyophilizable, and injectable.

The foregoing has been a description of certain preferred embodiments of the present invention, but is not intended to limit the invention in any way. Rather, many modifications, variations and changes in details may be made within the scope of the present invention.

What is claimed is:

- 1. A kit for forming a radiopharmaceutical imaging agent which includes a non-stannous reducing agent.
- 2. A kit according to claim 1, wherein said reducing agent is a metallic compound selected from the group consisting of Cu(I), Cu(II), Co(II), Fe(II), Sn(0), Zr(0), Cr(II) amd Zn(0).
- 3. A kit according to claim 1, wherein said reducing agent is a non-metallic compound selected from the group consisting of acids in general, dithionite, formamidine, formamadine sulfinic acid, phosphite, hypophosphite, dithiothreitol, hydrochloric acid, and borohydric acid.
- 4. A kit according to claim 1, wherein said reducing agent is selected from the group consisting of phosphines, sulfhydryl compounds, phosphites, thiols, thioethers, borates, borocyano groups, ascorbates, and gentisates.
- 5. A kit according to claim 4, wherein said reducing agent is a mono-dentate phosphine.
- 6. A kit according to claim 5, wherein said reducing agent is tris(3-methoxypropyl)phosphine.
- 7. A kit according to claim 5, wherein said reducing agent is a tertiary phosphine.
- 8. A kit according to claim 5, wherein said reducing agent is hypophosphite ion.

WO 94/22496 PCT/US94/03389

8

- 9. A kit according to claim 5, wherein said reducing agent is hydrogen phosphite.
- 10. A kit according to claim 1, wherein said reducing agent is also a complexing agent for the radiopharmaceutical imaging agent.

INTERNATIONAL SEARCH REPORT.

L. ./mational application No. PCT/US94/03389

			FC1/0394/033	
A. CLASSIFICATION OF SUBJECT MATTER				
IPC(5)	:A61K 49/02, 43/00 :424/1.11, 1.65			
	to International Patent Classification (IPC) or to both	national classification	and IPC	
	LDS SEARCHED			
Minimum o	documentation searched (classification system followe	d by classification sym	ibois)	
	424/1.11, 1.65	·, ·	,	
Documenta	tion searched other than minimum documentation to th	e extent that such docu	ments are included	in the fields searched
Electronic o	data base consulted during the international search (n	ame of data base and,	where practicable	, search terms used)
	ee Extra Sheet.	•	•	
C. DOC	CUMENTS CONSIDERED TO BE RELEVANT			· · · · · · · · · · · · · · · · · · ·
Category*	Citation of document, with indication, where a	nomoriste of the relev	ant page age	Relevant to claim No.
	The state of the s	ppropriate, or the relev	ant bessages	Rejevant to ciatri 140.
X	US, A, 3,749,556 (BARAK ET AL) 31 July 197	3. See the	1, 2
	Abstract; column 2, lines 47-61	; and column	3, line 68	·
	bridging column 4, line 3.			
V	110 4 4 200 200 444 54 54 54			
X	US, A, 4,208,398 (KUBIATOWIC	Z ET AL.) 17 J	lune 1980.	1-4, 8
Y	See column 6, lines 8-41.			0
•		•		9
X, P	US, A, 5,202,109 (FRITZBERG ET	AL.) 13 April	1993. See	1-4, 8
	column 16, lines 9-16 and column	19, lines 28-5	6.	·
x	US A 4 025 650 (NOCOC 57			
^	US, A, 4,925,650 (NOSCO ET / columns 5, lines 25-32 and lines (AL.) 15 May 1	990. See	1-3
	coloring 5, lines 25-32 and lines (00-00.		
x	US, A, 4,314,986 (RUDDOCK)	09 February 1	982. See	1. 2
1	column 2, lines 16-26.		, , ,	', -
X Further documents are listed in the continuation of Box C. See patent family annex.				
* Special extegories of cited documents: "T" Interdocument published after the international filing date or priority date and not in conflict with the application but cited to understand the				
"A" document defining the general state of the art which is not considered to be part of particular relevance				
E cartier document published on or after the international filling date "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step				
"L" document which may throw doubts on priority claim(s) or which is when the document is taken alone cited to establish the publication date of another citation or other				
special reason (as specified) "Y" document of particular relevance; the claimed invention cannot be considered to invention are inventive acts when the document is				
being obvious to a person skilled in the art				
*P" document published prior to the international filling date but later than *&* document member of the same patent family the priority date claimed				
Date of the actual completion of the international search Date of mailing of the international search report				
26 APRIL 1994 01 JUN 1994				
Name and mailing address of the ISA/US Authorized officer				
Commissioner of Patents and Trademarks Box PCT JOHN M. COVERT				
Washington Facsimile N	a, D.C. 20231 o. (703) 305-3230		03) 308-0444	
		1	,	

INTERNATIONAL SEARCH REPORT

i. .national application No. PCT/US94/03389

C (Continua	tion). DOCUMENTS CONSIDERED TO BE RELEVANT	
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 4,957,728 (DEUTSCH ET AL.) 18 September 1990. See column 2, line 1 bridging column 3, line 30; column 4, lines 1-15; and column 5, line 55 bridging column 6, line 6.	1, 2, 4-7, 10
x	US, A, 5,069,900 (LINDER) 03 DECEMBER 1991. See column 3, lines 45-65, and column 4, lines 37-58.	1, 4-7
A	US, A, 5,112,594 (WOULFE ET AL) 12 May 1992.	
	!	
:	,	

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

rnational application No. PCT/US94/03389

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS Messenger

L1 = 1427 S 424/1.1/CCLS

L2 = 9 S 424/1.!!/CCLS

L3 = 1436 L1 OR L2

L4 = 541 S ASCORB?/AB

L5 = 9 S L3 AND L4

L6 = 1007378 S REDUC?

L7 = 14740 S PHOSPHIT? OR HYPOPHOSPHIT?

L8 = 9 S L4 (P) L5 L9 = 18 S L7 AND L3

L10 = 113 S L4(P) L6

L11 = 1141 S TECHNETIUM OR PERTECHNETATE

L12 = 1 S L10(P) L11

L13 = 340244 S (REDUCING OR REDUCTANT)

L14 = 89 S L3 AND (L13 (P) (COPPER OR CU OR ZINC OR ZN OR IRON OR FE))

L15 = 7 S L 14 NOT L 9

L16 = 176 \$ 534/14/CCLS AND L3

LI7 = 17 S L16 AND PHOSPHIN?

L18 = 126 S L16 NOT (L14 OR L9)

L19 = 15 S L17 AND L13